Job Sheet for Seeding												
		50			ceang							
Planned by:	MO Mix 57 CP38A-33		Date:				County:					
Client:			Address:						<u> </u>			
Farm(s):		Tract(s):			Fields:			CIN#:		Ī		
Acres:	1.00	Program:								-		
Primary / Cost-shared FOTG Pra	actice:	Wildlife Habitat Plantir	ng (420)		Second	lary FOTG	Practice:					
Primary Purpose of Seeding:												
Site Soil Group:			OR	Ecol	ogical Site:							
Additional Information:												
Seedbed Preparation for Grass and/or Forb/Legume Seeding												
0 40 W	Seedbe	d Preparation	for Gra	ass an	d/or Foi	rb/Leg	ume S	eeding		_		
Current Conditions:										1		
	Row crop stubble or ti	lled seed bed (to inco	rporate ame	endments,	level field, c	or following	g construc	tion operations)				
	Killed sod (renovation;	for example from intr	oduced coc	ol season g	rasses to na	tive warm	season gr	asses or from warm s	season gra sses to			
Conditions Desired at Planting:	O Killed sod (renovation; for example from introduced cool season grasses to native warm season grasses or from warm season gra sses to a pollinator forb only planting)											
	O Suppressed existing vegetation, as for an inter-seeding of legumes on pastureland or native forbs on natural communities											
	Slopes that are 4:1 or s	O Slopes that are 4:1 or steeper that will receive a critical area planting (broadcast planting method only)										
Seed Mix Contains:	Cool Season Grasses	(majority)	/arm Seasor	n Grasses (	maiority)	Na	tive Forbs	Native/Intr	oduced Legumes	-		
Planting Method:		Interseeding:	No		ng Season:				outced Leguines			
Acceptable Planting Dates:		Optimal Plan			.g							
· · · · · · · · · · · · · · · · · · ·												
Seedbed Prep:												
Seed Placement:												
										-		
Additional Instructions: Fertilizer and Lime												
Planned												
Is Fertilizer and/or Lime Needed?	,											
Program rules and/or plann		re soil amendments (e	example - ge	eneral CRP	plantings, v	wildlife pla	ntings on	cropland)				
Program rules and/or plann						· · · ·		· ·	cent application of li	ming material		
and/or fertilizer). Apply fert			5									
Program rules and/or plann		nendments. Blanket ra	tes will be u	ised for a	Critical Area	Planting (	342) as inc	dicated below becaus	se a recent soil test i	<mark>s not availab</mark> le		
(as for constructed fills and	finished grades).											
Soil Test Recommendations			Limo*	N		KO						
Salt pH from soil test results:		lba/aa	Lime*	IN	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O						
		lbs/ac	Notrog	0	0	0						
		Total Needed	Not req.	0	0	0						
Displace action for a second state 1.00	uter e	lbs/ac	#N/A		<u>г г</u>							
Blanket rates for constructed fills when a lbs/ac #N/A current soil test is not available: Total Needed #N/A												
What legumes are included in the seed mix?												
When to apply lime:												
										-		
When to apply fertilizer:		Coil putricata are sti	hin merer	oouired (-	rtho plane -	d on c =!	thet	a plantad				
	No fertilizer will be applied.	Soli nutrients are wit	nin ranges i	equirea ta	i the planne	u species	unat Will De	e pianteo.		_ ,		

Additional Information:					
Companion Crop Specifications					
	hed prior to or at the same time as a permanent and managed to minimize competion with the p		r legumes.		
Planned Companion Crop:	None Planned Redtop Spring Oats	Winter Wheat	Rate per Acre (lbs):	Total Bulk Lbs to Seed:	
Additional Information:					
Applied					
Companion Crop Planted:	Oats Winter Wheat	Total L Bulk Lbs Planted:	bs Lbs/Ac		
			0.0		

## U.S. Department of Agriculture

## Natural Resources Conservation Service

Seeding Mix Summary												
Acres to be Seeded:	1.00											
Practice Adjustment Factor:			Broadca	st Adjustme	ent Factor:		_	Intersee	ding Adjus	tment Factor:	1	
Apply Adjustment Factors to:							-					
Planned Seed Mix				Seed Mix F	Purchased	and Plante	d					
							-					
Species	Acceptable Cultivar(s)	PLS Ibs/ac	Total PLS Ibs Needed	Bulk Ibs Seeded	% Purity	% Germ	% Dormant/ Hard Seed	PLS lbs Planted	Shortage/ Excess	% of Mix	Cultivar OR Origin for MO Source Identified Class Seed	Meets all Require- ments
Broomsedge (Bluestem), Andropogon virginicus	No cultivars available	0.440	0.440						-0.440			
Little Bluestem, Schizachyrium scoparium	Aldous (statewide), Camper (north MO), Cimarron (south MO - sandy sites only), Ozark (statewide), Yellow Tag (statewide)	0.730	0.730						-0.730			
Splitbeard Bluestem, Andropogon ternarius	No cultivars available	2.400	2.400						-2.400			
Black-eyed Susan (B), Rudbeckia		0.075	0.075						-0.075			
hirta Canada Milkvetch, Astragalus canadensis		0.020	0.020						-0.020			
Common Evening Primrose (B), Oenothera biennis		0.030	0.030						-0.030			
False Sunflower, Heliopsis helianthoides		0.050	0.050						-0.050			
Foxglove Beardtongue, Penstemon digitalis		0.019	0.019						-0.019			
Golden Alexanders, Zizia aurea		0.070	0.070						-0.070			
Gray Goldenrod, Solidago nemoralis		0.010	0.010						-0.010			
Gray-headed Coneflower, Ratibida pinnata	Available: Sunglow (statewide)	0.055	0.055						-0.055			
Illinois Bundleflower, Desmanthus illinoensis	Available: Reno Germplasm (statewide)	0.150	0.150						-0.150			
Lanceleaf Coreopsis, Coreopsis Ianceolata		0.100	0.100						-0.100			
Maximilian Sunflower, Helianthus maximiliani		0.010	0.010						-0.010			
Pale Purple Coneflower, Echinacea pallida		0.030	0.030						-0.030			

## U.S. Department of Agriculture Natural Resources Conservation Service

					-				-	-		
Partridge Pea (A), Chamaecrista fasciculata	Available: Riley (statewide), Lark Selection (statewide)	0.100	0.100						-0.100			
Prairie Cinquefoil, Drymocallis		0.035	0.035						-0.035			
arguta Purple Coneflower, Echinacea		0.135	0.135						-0.135			
purpurea		0.100	0.100						0.100			
Purple Prairie Clover, Dalea purpurea	Available: Kaneb (statewide)	0.128	0.128						-0.128			
Rattlesnake Master, Eryngium yuccifolium		0.020	0.020						-0.020			
Roundhead Lespedeza, Lespedeza capitata	Available: Kanoka (statewide)	0.020	0.020						-0.020			
Showy Ticktrefoil, Desmodium canadense	Available: Alexander Germplasm (statewide)	0.040	0.040						-0.040			
Smooth Aster, Symphyotrichum laeve		0.010	0.010						-0.010			
Stiff Goldenrod, Oligoneuron rigidum, Solidago rigida		0.010	0.010						-0.010			
Upright Prairie Coneflower, Ratibida columnifera		0.100	0.100						-0.100			
White Prairie Clover, Dalea candida		0.020	0.020						-0.020			
Wild Bergamot, Monarda fistulosa		0.070	0.070						-0.070			
	Total Planned:	4.877	4.877									
Planned Grasses PLS/ft <sup>2</sup>	Planned Grasses PLS/ft <sup>2</sup> : 24.01 Total PLS lbs. Forbs Planted:											
Planned Forbs/Legumes PLS/ft <sup>2</sup>	16.00	Total % of Mix:										
Planned Total PLS/ft <sup>2</sup>	Planned Total PLS/ft <sup>2</sup> : 40.01 % of mix of annual/biennial species:											
Customer and Seed Dealer Please Note:												

## **Customer and Seed Dealer Please Note:**

Only viable, high quality and adapted seed will be used. All seed shall have a current seed test within 10 months of the planting date that lists germination, purity, and hard seed as a percentage for determining pure live seed and lists the percent of weed seed present that meets State seed quality law standards. Seed test results must be provided to the customer and USDA for confirmation that seed planted meets specifications outlined in this plan. Seed must be clean and relatively free of weed seed and contaminants. Seed that has become wet, moldy, or otherwise damaged in transit or storage is not acceptable. Certified seed is preferred but not required unless Missouri Source Identified Class Seed, or yellow tag, is denoted as being required.

Legume seed shall be inoculated with the proper, viable Rhizobium bacteria species prior to planting according to Missouri Agronomy Technical Note MO-36, "Legume Inoculation." Pre-inoculated seed shall be planted prior to the expiration date on the inoculum tag or be re-inoculated with the appropriate inoculum within 24 hours prior to seeding. Inoculated seed will not be exposed to direct sunlight. Inoculation of legumes may be waived when a current pasture inventory (JS-AGRON-24 or similar inventory) shows that the legume species to be seeded occupies more than 5 percent of the vegetative cover on the site prior to seeding.

For "Acceptable Cultivar(s)," north Missouri (MO) is all counties north of Bates, Henry, Benton, Morgan, Moniteau, Cole, Osage, Gasconade, Franklin, and St. Louis Counties. South Missouri is all counties including and south of those listed. Refer to Native Cultivars or Selections Information Sheet (IS-MO723N) for guidance on selecting the appropriate native grass cultivars to meet objectives. In all cases where a cultivar(s) is/are noted under the Acceptable Cultivar(s) heading above, Missouri Source Identified Class Seed (yellow tag seed) may be used instead of a cultivar variety or selection.

The following information applies only if "Locally adapted Missouri Source Identified Class Seed (yellow tag) required" is shown under the "Acceptable Cultivar(s)" heading above: Locally adapted Missouri Source Identified Class Seed, or yellow tag, is defined as a native plant that source genetically originated in Missouri, was not introduced, and existed within the state borders prior to arrival of settlers. All seed from herbaceous material shall comply with Missouri seed laws including Missouri Crop Improvement Association guidance. All seed will comply with AOSCA (Association of Official Seed Certifying Agencies) certification procedures (including appropriate tagging) to include third-party verification by the Missouri origin (grown in Missouri) and certified as Missouri Source Identified Class. Source Identified Class. Source Identified Class. Source Identified Cretification means: parent seed is collected from natural remnant Missouri populations; no selection, testing, or breeding for specific traits; production fields are inspected to verify species, source, and lack of noxious weeds; and seed is certified for purity and germination. If Missouri origin (grown) source identified class seed is not available, Missouri source identified class seed may be obtained from adjoining states.

ľ

1

Mulching Specifications (After Grass Seeding)									
Type of Mulch Required: Straw, Grass Hay, Alfalfa Hay - 1 1/2 to 2 1/2 Tons/Acre									
Amount of Mulch Required for This Site:									
Minimum Rate (lbs):	3,000 OR	Tons: 1.50	Maximum Rate (lbs): 5,000	OR Tons:	2.5				
Additional Information: n/a									
Applied									
Type of Mulch Used:									
Number of Bales:	Bale	Veight, lbs:	Mulch Applied, Tons:	Excess/Shor	tage, Tons:				
Based on the information in this job shee	Based on the information in this job sheet, all requirements for this seeding should be completed by:								
Was site prep completed as	s outlined above?								
Seeding was completed on		to the above requirements							
	(Date)								
Producer's or Contr	actor's Signature		Date						
For NRCS Use Only:									
Does this seeding meet all NRCS standard	Does this seeding meet all NRCS standards and specifications and was implemented as planned above?								
Certified by:									
NRCS Representative Date Field Office									
Additional Comments:									
When seeding is completed, return seeding plan to the Natural Resources Conservation Service with all receipts pertaining to this seeding (seed, fertilzer, lime, mulch,									
companion crop, labor, drill rental, etc.) and seed tests, including tags for yellow tag seed.									