| Job Sheet for Seeding                                 |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|---|--|---|---------------|-------------|---------------|------------------|--------------|----------------------|---------------------------------|-------------------------------|--|--|--|
|   |  |   |               |             | eeuing        |                  |              |                      |                                 |                               |  |  |  |
| Planned by:   | MO Mix 59 CP38A-43   |   | Date:         |             |               |                  | County:      |                      |                                 |                               |  |  |  |
| Client:   |  |   | Address:      |             |               |                  | countyr      |                      | <u> </u>                        |                               |  |  |  |
| Farm(s):  |  | Tract(s):   | / laar ooo.   |             | Fields:       |                  |              | CIN#:                |                                 | -                             |  |  |  |
| Acres:  | 1.00   | Program:  |               |             | T Ioldo.      |                  |              |                      | <u> </u>                        | 4                             |  |  |  |
| Primary / Cost-shared FOTG Pra                        |  | Wildlife Habitat Plantir  | ng (420)      |             | Second        | lary FOTG        | Practice:    |                      |                                 |                               |  |  |  |
| Primary Purpose of Seeding:                           |  |   | .9 (120)      |             |               |                  | T Tuotico.   |                      |                                 |                               |  |  |  |
| Site Soil Group:                                      |  |   | OR            | Ecol        | ogical Site:  |                  |              |                      |                                 |                               |  |  |  |
| Additional Information:                               |  |   | on            | 2001        | oglour onto.  |                  |              |                      |                                 | -                             |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 | <u>_</u>                      |  |  |  |
|   | Seedbe   | d Preparation   | for Gra       | ass an      | d/or Foi      | rb/Lea           | ume S        | eedina               |                                 |                               |  |  |  |
| Current Conditions:                                   |  |   |               |             |               |                  |              | <u> </u>             |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 | 1                             |  |  |  |
|   | Row crop stubble or ti   |   |               |             |               |                  |              | •                    |                                 |                               |  |  |  |
|   | Killed sod (renovation;  | for example from intr   | oduced coc    | ol season g | rasses to na  | tive warm        | season gr    | asses or from warm s | <mark>season gra sses</mark> to |                               |  |  |  |
| Conditions Desired at Planting:                       | a pollinator forb only planting)   |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   | O Suppressed existing vegetation, as for an inter-seeding of legumes on pastureland or native forbs on natural communities |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   | Slopes that are 4:1 or s   | at are 4:1 or steeper that will receive a critical area planting (broadcast planting method only) |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               | -           |               |                  |              |                      |                                 | -                             |  |  |  |
| Seed Mix Contains:                                    | Cool Season Grasses  |   | /arm Seasor   |             |               | L Na             | tive Forbs   | Native/Intr          | oduced Legumes                  |                               |  |  |  |
| Planting Method:                                      |  | Interseeding:   | No            | Plantir     | ng Season:    |                  |              |                      |                                 |                               |  |  |  |
| Acceptable Planting Dates:                            |  | Optimal Plan  | ting Dates:   |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Seedbed Prep:   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 | -                             |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Seed Placement:                                       |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Additional Instructions:                              |  |   |               |             |               |                  |              |                      |                                 | -                             |  |  |  |
| Fertilizer and Lime                                   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Planned   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Is Fertilizer and/or Lime Needed?                     |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Program rules and/or plann                            |  | re soil amendments (e   | example - ge  | eneral CRP  | plantings, v  | vildlife pla     | ntinas on    | cropland)            |                                 |                               |  |  |  |
| Program rules and/or plann                            |  |   |               |             |               | · · · ·          |              |                      | cent application of li          | ming material                 |  |  |  |
| and/or fertilizer). Apply fert                        |  |   | 5             |             |               | 55 than 4 y      |              |                      |                                 | ming material                 |  |  |  |
| Program rules and/or plann                            |  | nendments. Blanket ra   | tes will be u | ised for a  | Critical Area | Planting (       | 342) as inc  | dicated below becaus | se a recent soil test i         | <mark>s not availabl</mark> e |  |  |  |
| (as for constructed fills and                         | finished grades).  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Soil Test Recommendations                             |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Salt pH from soil test results:                       |  |   | Lime*         | Ν           | $P_2O_5$      | K <sub>2</sub> O |              |                      |                                 |                               |  |  |  |
|   |  | lbs/ac  |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  | Total Needed  | Not req.      | 0           | 0             | 0                |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| Blanket rates for constructed fills                   | when a   | lbs/ac  | #N/A          |             |               |                  |              |                      |                                 |                               |  |  |  |
| current soil test is not available: Total Needed #N/A |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| What legumes are included in the                      | e seed mix?  | e Alfalfa   | Cool Season   | Legumes     | other than /  | Alfalfa          | Wa           | rm Season Legumes    |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
| When to apply lime:                                   |  |   |               |             |               |                  |              |                      |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 | -                             |  |  |  |
| When to apply fertilizer:                             | No fertilizer will be applied.   | Soil nutrients are wit  | hin ranges r  | reauired fo | r the planne  | d species        | that will be | e planted.           |                                 |                               |  |  |  |
|   |  |   |               |             |               |                  |              |                      |                                 | -                             |  |  |  |

| Additional Information:       |  |                              |                      |                         |  |
|-------------------------------|--|------------------------------|----------------------|-------------------------|--|
|                               |  |                              |                      |                         |  |
|                               |  |                              |                      |                         |  |
| Companion Crop Specifications |  |                              |                      |                         |  |
|                               | hed prior to or at the same time as a permanent and managed to minimize competion with the p |                              | r legumes.           |                         |  |
| Planned Companion Crop:       | None Planned Redtop Spring Oats  | Winter Wheat                 | Rate per Acre (lbs): | Total Bulk Lbs to Seed: |  |
|                               |  |                              |                      |                         |  |
| Additional Information:       |  |                              |                      |                         |  |
|                               |  |                              |                      |                         |  |
|                               |  |                              |                      |                         |  |
| Applied                       |  |                              |                      |                         |  |
| Companion Crop Planted:       | Oats Winter Wheat  | Total L<br>Bulk Lbs Planted: | bs Lbs/Ac            |                         |  |
|                               |  |                              | 0.0                  |                         |  |

## U.S. Department of Agriculture

## Natural Resources Conservation Service

| Seeding Mix Summary                               |  |               |                         |                    |             |        |                               |                    |                     |               |  |                                   |
|---|--|---------------|-------------------------|--------------------|-------------|--------|-------------------------------|--------------------|---------------------|---------------|--|-----------------------------------|
| Acres to be Seeded:                               | 1.00   |               |                         |                    |             |        |                               |                    |                     |               |  |                                   |
| Practice Adjustment Factor:                       |  |               | Broadca                 | st Adjustme        | ent Factor: |        |                               | Intersee           | ding Adjus          | tment Factor: | 1  |                                   |
| Apply Adjustment Factors to:                      |  |               |                         |                    |             |        |                               |                    |                     |               |  |                                   |
| Planned Seed Mix Seed Mix Purchased and Planted   |  |               |                         |                    |             |        |                               |                    |                     |               |  |                                   |
|   |  |               |                         |                    |             |        |                               |                    |                     |               |  |                                   |
| Species   | Acceptable Cultivar(s)   | PLS<br>Ibs/ac | Total PLS<br>Ibs Needed | Bulk lbs<br>Seeded | % Purity    | % Germ | %<br>Dormant/<br>Hard<br>Seed | PLS lbs<br>Planted | Shortage/<br>Excess | % of Mix      | Cultivar OR<br>Origin<br>for MO Source<br>Identified Class<br>Seed | Meets<br>all<br>Require-<br>ments |
| Broomsedge (Bluestem),<br>Andropogon virginicus   | No cultivars available   | 0.280         | 0.280                   |                    |             |        |                               |                    | -0.280              |               |  |                                   |
| Little Bluestem, Schizachyrium<br>scoparium       | Aldous (statewide), Camper<br>(north MO), Cimarron (south<br>MO - sandy sites only),<br>Ozark (statewide), Yellow<br>Tag (statewide) | 1.350         | 1.350                   |                    |             |        |                               |                    | -1.350              |               |  |                                   |
| Splitbeard Bluestem, Andropogon<br>ternarius      | No cultivars available   | 2.500         | 2.500                   |                    |             |        |                               |                    | -2.500              |               |  |                                   |
| Black-eyed Susan (B), Rudbeckia<br>hirta          |  | 0.110         | 0.110                   |                    |             |        |                               |                    | -0.110              |               |  |                                   |
| Blue Lobelia, Lobelia siphilitica                 |  | 0.008         | 0.008                   |                    |             |        |                               |                    | -0.008              |               |  |                                   |
| Canada Milkvetch, Astragalus<br>canadensis        |  | 0.076         | 0.076                   |                    |             |        |                               |                    | -0.076              |               |  |                                   |
| Common Evening Primrose (B),<br>Oenothera biennis |  | 0.020         | 0.020                   |                    |             |        |                               |                    | -0.020              |               |  |                                   |
| Common Milkweed, Asclepias<br>syriaca             |  | 0.510         | 0.510                   |                    |             |        |                               |                    | -0.510              |               |  |                                   |
| False Sunflower, Heliopsis<br>helianthoides       |  | 0.140         | 0.140                   |                    |             |        |                               |                    | -0.140              |               |  |                                   |
| Foxglove Beardtongue,<br>Penstemon digitalis      |  | 0.013         | 0.013                   |                    |             |        |                               |                    | -0.013              |               |  |                                   |

| Oenothera biennis                                |  | 0.020 | 0.020 |  |  | -0.020 |  |  |
|--|--|-------|-------|--|--|--------|--|--|
| Common Milkweed, Asclepias<br>syriaca            |  | 0.510 | 0.510 |  |  | -0.510 |  |  |
| False Sunflower, Heliopsis<br>helianthoides      |  | 0.140 | 0.140 |  |  | -0.140 |  |  |
| Foxglove Beardtongue,<br>Penstemon digitalis     |  | 0.013 | 0.013 |  |  | -0.013 |  |  |
| Golden Alexanders, Zizia aurea                   |  | 0.090 | 0.090 |  |  | -0.090 |  |  |
| Gray Goldenrod, Solidago<br>nemoralis            |  | 0.012 | 0.012 |  |  | -0.012 |  |  |
| Gray-headed Coneflower, Ratibida<br>pinnata      | Available: Sunglow<br>(statewide)        | 0.050 | 0.050 |  |  | -0.050 |  |  |
| Illinois Bundleflower, Desmanthus<br>illinoensis | Available: Reno Germplasm<br>(statewide) | 0.150 | 0.150 |  |  | -0.150 |  |  |
| Lanceleaf Coreopsis, Coreopsis<br>lanceolata     |  | 0.200 | 0.200 |  |  | -0.200 |  |  |
| Maximilian Sunflower, Helianthus maximiliani     |  | 0.010 | 0.010 |  |  | -0.010 |  |  |
| Pale Purple Coneflower,<br>Echinacea pallida     |  | 0.050 | 0.050 |  |  | -0.050 |  |  |

## U.S. Department of Agriculture Natural Resources Conservation Service

| Partridge Pea (A), Chamaecrista<br>fasciculata           | Available: Riley (statewide),<br>Lark Selection (statewide)                    | 0.130           | 0.130 |  |           |             |          |  | -0.130 |  |  |  |
|--|--|-----------------|-------|--|-----------|-------------|----------|--|--------|--|--|--|
| Prairie Cinquefoil, Drymocallis<br>arguta                |  | 0.048           | 0.048 |  |           |             |          |  | -0.048 |  |  |  |
| Purple Coneflower, Echinacea<br>purpurea                 |  | 0.150           | 0.150 |  |           |             |          |  | -0.150 |  |  |  |
| Purple Prairie Clover, Dalea<br>purpurea                 | Available: Kaneb (statewide)   | 0.254           | 0.254 |  |           |             |          |  | -0.254 |  |  |  |
| Rattlesnake Master, Eryngium<br>yuccifolium              |  | 0.020           | 0.020 |  |           |             |          |  | -0.020 |  |  |  |
| Roundhead Lespedeza,<br>Lespedeza capitata               | Available: Kanoka<br>(statewide)   | 0.020           | 0.020 |  |           |             |          |  | -0.020 |  |  |  |
| Showy Ticktrefoil, Desmodium<br>canadense                | Available: Alexander<br>Germplasm (statewide)                                  | 0.040           | 0.040 |  |           |             |          |  | -0.040 |  |  |  |
| Smooth Aster, Symphyotrichum<br>laeve                    |  | 0.003           | 0.003 |  |           |             |          |  | -0.003 |  |  |  |
| Stiff Goldenrod, Oligoneuron<br>rigidum, Solidago rigida |  | 0.010           | 0.010 |  |           |             |          |  | -0.010 |  |  |  |
| Upright Prairie Coneflower,<br>Ratibida columnifera      |  | 0.115           | 0.115 |  |           |             |          |  | -0.115 |  |  |  |
| White Prairie Clover, Dalea<br>candida                   |  | 0.098           | 0.098 |  |           |             |          |  | -0.098 |  |  |  |
| Wild Bergamot, Monarda fistulosa                         |  | 0.120           | 0.120 |  |           |             |          |  | -0.120 |  |  |  |
| Yarrow, Achillea millefolium                             |  | 0.009           | 0.009 |  |           |             |          |  | -0.009 |  |  |  |
|  |  |                 |       |  |           |             |          |  |        |  |  |  |
|  | Total Planned:   | 6.586           | 6.586 |  |           |             |          |  |        |  |  |  |
| Planned Grasses PLS/ft <sup>2</sup> :                    | 25.00  |                 |       |  | Total PLS | Blbs. Forbs | Planted: |  |        |  |  |  |
| Planned Forbs/Legumes PLS/ft <sup>2</sup> :              | 25.00  | Total % of Mix: |       |  |           |             |          |  |        |  |  |  |
| Planned Total PLS/ft <sup>2</sup> :                      | Planned Total PLS/ft <sup>2</sup> : 50.00 % of mix of annual/biennial species: |                 |       |  |           |             |          |  |        |  |  |  |
| Customer and Seed Dealer Please                          | e Note:  |                 |       |  |           |             |          |  |        |  |  |  |

Customer and Seed Dealer Please Note:

## U.S. Department of Agriculture Natural Resources Conservation Service

Only viable, high quality and adapted seed will be used. All seed shall have a current seed test within 10 months of the planting date that lists germination, purity, and hard seed as a percentage for determining pure live seed and lists the percent of weed seed present that meets State seed quality law standards. Seed test results must be provided to the customer and USDA for confirmation that seed planted meets specifications outlined in this plan. Seed must be clean and relatively free of weed seed and contaminants. Seed that has become wet, moldy, or otherwise damaged in transit or storage is not acceptable. Certified seed is preferred but not required unless Missouri Source Identified Class Seed, or yellow tag, is denoted as being required.

Legume seed shall be inoculated with the proper, viable Rhizobium bacteria species prior to planting according to Missouri Agronomy Technical Note MO-36, "Legume Inoculation." Pre-inoculated seed shall be planted prior to the expiration date on the inoculum tag or be re-inoculated with the appropriate inoculum within 24 hours prior to seeding. Inoculated seed will not be exposed to direct sunlight. Inoculation of legumes may be waived when a current pasture inventory (JS-AGRON-24 or similar inventory) shows that the legume species to be seeded occupies more than 5 percent of the vegetative cover on the site prior to seeding.

For "Acceptable Cultivar(s)," north Missouri (MO) is all counties north of Bates, Henry, Benton, Morgan, Moniteau, Cole, Osage, Gasconade, Franklin, and St. Louis Counties. South Missouri is all counties including and south of those listed. Refer to Native Cultivars or Selections Information Sheet (IS-MO723N) for guidance on selecting the appropriate native grass cultivars to meet objectives. In all cases where a cultivar(s) is/are noted under the Acceptable Cultivar(s) heading above, Missouri Source Identified Class Seed (yellow tag seed) may be used instead of a cultivar variety or selection.

The following information applies only if "Locally adapted Missouri Source Identified Class Seed (yellow tag) required" is shown under the "Acceptable Cultivar(s)" heading above: Locally adapted Missouri Source Identified Class Seed, or yellow tag, is defined as a native plant that source genetically originated in Missouri, was not introduced, and existed within the state borders prior to arrival of settlers. All seed from herbaceous material shall comply with Missouri seed laws including Missouri Crop Improvement Association guidance. All seed will comply with AOSCA (Association of Official Seed Certifying Agencies) certification procedures (including appropriate tagging) to include third-party verification by the Missouri Crop Improvement Association of source, genetic identity, and genetic purity of wildland collected or field or nursery grown plant germplasm materials. Seed must be Missouri origin (grown in Missouri) and certified as Missouri Source Identified Class. Source Identified Class. Source Identified Certification means: parent seed is collected from natural remnant Missouri populations; no selection, testing, or breeding for specific traits; production fields are inspected to verify species, source, and lack of noxious weeds; and seed is certified for purity and germination. If Missouri origin (grown) source identified class seed is not available, Missouri source identified class seed may be obtained from adjoining states.

| Mulching Specifications (After Grass Seeding)  |  |  |                           |              |             |  |  |  |  |  |
|--|--|--|---------------------------|--------------|-------------|--|--|--|--|--|
| Type of Mulch Required: None   | Straw, Grass Hay, A  | lfalfa Hay - 1 1/2 to 2 1/2 <sup>-</sup> | Tons/Acre                 |              |             |  |  |  |  |  |
| Amount of Mulch Required for This Site:  |  |  |                           |              |             |  |  |  |  |  |
| Minimum Rate (lbs):  | 3,000 OF   | R Tons: 1.50                             | Maximum Rate (lbs): 5,000 | OR Tons:     | 2.5         |  |  |  |  |  |
|  |  |  |                           |              |             |  |  |  |  |  |
| Additional Information: n/a  |  |  |                           |              |             |  |  |  |  |  |
|  |  |  |                           |              |             |  |  |  |  |  |
|  |  |  |                           |              |             |  |  |  |  |  |
| Applied  |  |  |                           |              |             |  |  |  |  |  |
| Type of Mulch Used:  |  |  |                           |              |             |  |  |  |  |  |
| Number of Bales:   | Bale   | Weight, lbs:                             | Mulch Applied, Tons:      | Excess/Shor  | tage, Tons: |  |  |  |  |  |
|  |  |  |                           |              |             |  |  |  |  |  |
| Based on the information in this job shee  | et, all requirements for this s  | eeding should be complete                | ed by:                    | Year:        |             |  |  |  |  |  |
| Was site prep completed as   | s outlined above?  |  |                           |              |             |  |  |  |  |  |
| Seeding was completed on   |  | g to the above requiremer                | its.                      |              |             |  |  |  |  |  |
|  | (Date)   |  |                           |              |             |  |  |  |  |  |
|  |  |  |                           |              |             |  |  |  |  |  |
| Producer's or Contr  | actor's Signature  |  | Date                      |              |             |  |  |  |  |  |
| For NRCS Use Only:   | For NRCS Use Only:   |  |                           |              |             |  |  |  |  |  |
| Does this seeding meet all NRCS standard   | Does this seeding meet all NRCS standards and specifications and was implemented as planned above? |  |                           |              |             |  |  |  |  |  |
| Certified by:  |  |  |                           |              |             |  |  |  |  |  |
| NRCS   | Representative   |  | Date                      | Field Office |             |  |  |  |  |  |
| Additional Comments:   |  |  |                           |              |             |  |  |  |  |  |
| When seeding is completed, return seeding plan to the Natural Resources Conservation Service with all receipts pertaining to this seeding (seed, fertilzer, lime, mulch, |  |  |                           |              |             |  |  |  |  |  |
| companion crop, labor, drill rental, etc.) and seed tests, including tags for yellow tag seed.   |  |  |                           |              |             |  |  |  |  |  |
|  |  |  |                           |              |             |  |  |  |  |  |