| | | Jo | b Sheet for Se | eding | | | | |
|---|---|--|--|--|----------------------|-------------------------|----------------|--|
| | | | | | | | | |
| Planned by: | MO Mix 23 - CP38E-4D | | Date: | | County: | | | |
| Client: | | T (1) | Address: | E: 11 | | O.N.I. | | |
| Farm(s): | 4.00 | Tract(s): | | Fields: | | CIN#: | | |
| Acres: | 1.00 | Program: | a (420) | Casardan, FO | TC Dresties | | | |
| Primary / Cost-shared FOTG Pra | actice: | Wildlife Habitat Planting | g (420) | Secondary FO | TG Practice: | | | |
| Primary Purpose of Seeding: Site Soil Group: | | | OR Ecolo | gical Site: | | | | |
| Additional Information: | | | OK ECOIO | gical Site. | | | | |
| Additional information. | | | | | | | | |
| | Seedbe | ed Preparation | for Grass and | /or Forb/Le | egume See | ding | | |
| Current Conditions: | | | | | | | | |
| Conditions Desired at Planting: | Row crop stubble or to Killed sod (renovation a pollinator forb only) Suppressed existing v Slopes that are 4:1 or | n; for example from intro planting) regetation, as for an inte | oduced cool season gr | asses to native war | r native forbs on | es or from warm seasc | on gra sses to | |
| Seed Mix Contains: | Cool Season Grasse | s (majority) W: | arm Season Grasses (r | naiority) | Native Forbs | Native/Introduc | red Legumes | |
| Planting Method: | Cool Season Glasse | Interseeding: | | Season: | INALIVE I OIDS | Native/introduc | led Legames | |
| Acceptable Planting Dates: | | Optimal Plant | | | | | | |
| | | | | | | | | |
| Seedbed Prep: | | | | | | | | |
| Seed Placement: | | | | | | | | |
| Additional Instructions: | | | | | | | | |
| Fertilizer and Lime | | | | | | | | |
| Planned | | | | | | | | |
| Is Fertilizer and/or Lime Needed? | | | | | | | | |
| Program rules and/or plann Program rules and/or plann and/or fertilizer). Apply fert Program rules and/or plann (as for constructed fills and | ed practices require soil a ilizer and/or lime accordin ed practices require soil a | mendments be applied a | according to a recent adations as indicated b | soil test (less than pelow. | 4 years old and a | after the most recent a | | |
| Soil Test Recommendations | | | | | | | | |
| Salt pH from soil test results: | | Γ | Lime* N | P ₂ O ₅ K ₂ O | | | | |
| | | lbs/ac Total Needed | Not req. 0 | 0 0 | | | | |
| Blanket rates for constructed fills current soil test is not available | | lbs/ac Total Needed | #N/A #N/A | | 7 | | | |
| What legumes are included in the | e seed mix? | ne Alfalfa C | ool Season Legumes o | ther than Alfalfa | Warm S | Season Legumes | | |
| When to apply lime: | | | | | | | | |
| When to apply fertilizer: | No fertilizer will be applied | Soil nutrients are with | in ranges required for | the planned speci | ies that will be pla | anted. | <u> </u> | |

U.S. Department of Agriculture Natural Resources Conservation Service

Revised 9/2017

| Additional Information: | | | | | | |
|---|---------------|--------------------|-------------------|----------------------|-------------------------|--|
| | | | | | | |
| | | | | | _ | |
| | | | | | | |
| | | | | | | |
| | - | | | | | |
| | | | | | | |
| Companion Crop Specifications | ; | | | | | |
| A companion crop may be estable The companion crop will be selec | | | | d/or legumes. | | |
| Planned Companion Crop: | None Planned | Redtop Spring Oats | Winter Wheat | Rate per Acre (lbs): | Total Bulk Lbs to Seed: | |
| | | | | | _ | |
| Additional Information: | | | | | | |
| | | | | | | |
| | | | | | | |
| Applied | | | | | | |
| | | | | | | |
| Companion Crop Planted: | | | Tot | tal Lbs Lbs/Ac | | |
| Redtop Spring | g Oats Winter | r Wheat | Bulk Lbs Planted: | 0.0 | | |

| Seeding Mix Summary | | | | | | | | | | | | |
|---|--|---------------|-------------------------|--------------------|-------------|--------|-------------------------------|--------------------|---------------------|---------------|--|-----------------------------------|
| Acres to be Seeded: | 1.00 | | | | | | | | | | | |
| Practice Adjustment Factor: | | • | Broadca | st Adjustme | ent Factor: | | - | Intersee | ding Adjus | tment Factor: | 1 | <u>-</u> |
| Apply Adjustment Factors to: | | | | | | | | | | | | |
| Planned Seed Mix Seed Mix Purchased and Planted | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| Species | Acceptable Cultivar(s) | PLS lbs/ac | Total PLS lbs Needed | Bulk lbs Seeded | % Purity | % Germ | % Dormant/ Hard Seed | PLS lbs Planted | Shortage/ Excess | % of Mix | Cultivar OR Origin for MO Source Identified Class Seed | Meets all Require- ments |
| Little Bluestem, Schizachyrium scoparium | Aldous (statewide), Camper (north MO), Cimarron (south MO - sandy sites only), Ozark (statewide), Yellow Tag (statewide) | 1.086 | 1.086 | | | | | | -1.086 | | | |
| Sideoats Grama, Bouteloua curtipendula | Butte (north MO), El Reno (statewide), Trailway (statewide), Yellow Tag (statewide) | 1.094 | 1.094 | | | | | | -1.094 | | | |
| Black-eyed Susan (B), Rudbeckia hirta | | 0.030 | 0.030 | | | | | | -0.030 | | | |
| Blue Lobelia, Lobelia siphilitica | | 0.005 | 0.005 | | | | | | -0.005 | | | |
| Canada Milkvetch, Astragalus | | 0.250 | 0.250 | | | | | | -0.250 | | | |
| canadensis Common Evening Primrose (B), | | 0.030 | 0.030 | | | | | | -0.030 | | | |
| Oenothera biennis Common Milkweed, Asclepias | | | | | | | | | | | | |
| syriaca False Sunflower, Heliopsis | | 0.600 | 0.600 | | | | | | -0.600 | | | |
| helianthoides | | 0.400 | 0.400 | | | | | | -0.400 | | | |
| Foxglove Beardtongue, Penstemon digitalis | | 0.010 | 0.010 | | | | | | -0.010 | | | |
| Golden Alexanders, Zizia aurea | | 0.250 | 0.250 | | | | | | -0.250 | | | |
| Gray Goldenrod, Solidago nemoralis | | 0.140 | 0.140 | | | | | | -0.140 | | | |
| Gray-headed Coneflower, Ratibida pinnata | Available: Sunglow (statewide) | 0.150 | 0.150 | | | | | | -0.150 | | | |
| Illinois Bundleflower, Desmanthus illinoensis | Available: Reno Germplasm (statewide) | 0.320 | 0.320 | | | | | | -0.320 | | | |
| Lanceleaf Coreopsis, Coreopsis lanceolata | | 0.250 | 0.250 | | | | | | -0.250 | | | |
| Maximilian Sunflower, Helianthus | | 0.250 | 0.250 | | | | | | -0.250 | | | |
| maximiliani | | | -: | | | | | | | | | |

| Pale Purple Coneflower, | | | | | | | | | | | |
|--|---|-------|-------|--|-----------|--------------|----------|---|--------|--|--|
| Echinacea pallida | | 0.190 | 0.190 | | | | | | -0.190 | | |
| Partridge Pea (A), Chamaecrista fasciculata | Available: Riley (statewide), Lark Selection (statewide) | 0.250 | 0.250 | | | | | | -0.250 | | |
| Prairie Cinquefoil, Drymocallis arguta | | 0.010 | 0.010 | | | | | | -0.010 | | |
| Purple Prairie Clover, Dalea purpurea | Available: Kaneb (statewide) | 0.250 | 0.250 | | | | | | -0.250 | | |
| Rattlesnake Master, Eryngium yuccifolium | | 0.020 | 0.020 | | | | | | -0.020 | | |
| Roundhead Lespedeza, Lespedeza capitata | Available: Kanoka (statewide) | 0.100 | 0.100 | | | | | | -0.100 | | |
| Showy Ticktrefoil, Desmodium canadense | Available: Alexander Germplasm (statewide) | 0.120 | 0.120 | | | | | | -0.120 | | |
| Smooth Aster, Symphyotrichum laeve | | 0.010 | 0.010 | | | | | | -0.010 | | |
| Stiff Goldenrod, Oligoneuron rigidum, Solidago rigida | | 0.100 | 0.100 | | | | | | -0.100 | | |
| Swamp Milkweed, Asclepias incarnata | | 0.150 | 0.150 | | | | | | -0.150 | | |
| Upright Prairie Coneflower, Ratibida columnifera | | 0.160 | 0.160 | | | | | | -0.160 | | |
| White Prairie Clover, Dalea candida | | 0.400 | 0.400 | | | | | | -0.400 | | |
| Wild Bergamot, Monarda fistulosa | | 0.050 | 0.050 | | | | | | -0.050 | | |
| Yarrow, Achillea millefolium | | 0.009 | 0.009 | | | | | | -0.009 | | |
| | | | | | | | | | | | |
| | Total Planned: | 6.684 | 6.684 | | | | | | | | |
| Planned Grasses PLS/ft ² : | 10.00 | | | | Total PLS | S lbs. Forbs | Planted: | - | | | |
| Planned Forbs/Legumes PLS/ft ² : 30.00 Total % of Mix: | | | | | | | | | | | |
| Planned Total PLS/ft ² : 40.00 % of mix of annual/biennial species: | | | | | | | | | | | |
| Customer and Seed Dealer Please Note: | | | | | | | | | | | |

Only viable, high quality and adapted seed will be used. All seed shall have a current seed test within 10 months of the planting date that lists germination, purity, and hard seed as a percentage for determining pure live seed and lists the percent of weed seed present that meets State seed quality law standards. Seed test results must be provided to the customer and USDA for confirmation that seed planted meets specifications outlined in this plan. Seed must be clean and relatively free of weed seed and contaminants. Seed that has become wet, moldy, or otherwise damaged in transit or storage is not acceptable. Certified seed is preferred but not required unless Missouri Source Identified Class Seed, or yellow tag, is denoted as being required.

Legume seed shall be inoculated with the proper, viable Rhizobium bacteria species prior to planting according to Missouri Agronomy Technical Note MO-36, "Legume Inoculation." Pre-inoculated seed shall be planted prior to the expiration date on the inoculum tag or be re-inoculated with the appropriate inoculum within 24 hours prior to seeding. Inoculated seed will not be exposed to direct sunlight. Inoculation of legumes may be waived when a current pasture inventory (JS-AGRON-24 or similar inventory) shows that the legume species to be seeded occupies more than 5 percent of the vegetative cover on the site prior to seeding.

For "Acceptable Cultivar(s)," north Missouri (MO) is all counties north of Bates, Henry, Benton, Morgan, Moniteau, Cole, Osage, Gasconade, Franklin, and St. Louis Counties. South Missouri is all counties including and south of those listed. Refer to Native Cultivars or Selections Information Sheet (IS-MO723N) for guidance on selecting the appropriate native grass cultivars to meet objectives. In all cases where a cultivar(s) is/are noted under the Acceptable Cultivar(s) heading above, Missouri Source Identified Class Seed (yellow tag seed) may be used instead of a cultivar variety or selection.

The following information applies only if "Locally adapted Missouri Source Identified Class Seed (yellow tag) required" is shown under the "Acceptable Cultivar(s)" heading above: Locally adapted Missouri Source Identified Class Seed, or yellow tag, is defined as a native plant that source genetically originated in Missouri, was not introduced, and existed within the state borders prior to arrival of settlers. All seed from herbaceous material shall comply with Missouri seed laws including Missouri Crop Improvement Association guidance. All seed will comply with AOSCA (Association of Official Seed Certifying Agencies) certification procedures (including appropriate tagging) to include third-party verification by the Missouri Crop Improvement Association of source, genetic identity, and genetic purity of wildland collected or field or nursery grown plant germplasm materials. Seed must be Missouri origin (grown in Missouri) and certified as Missouri Source Identified Class. Source Identified Certification means: parent seed is collected from natural remnant Missouri populations; no selection, testing, or breeding for specific traits; production fields are inspected to verify species, source, and lack of noxious weeds; and seed is certified for purity and germination. If Missouri origin (grown) source identified class seed is not available, Missouri source identified class seed may be obtained from adjoining states.

| Mulching Specifications (After Grass Seeding) | | | | | | | | | | |
|---|--|-------------------------|---------------------|-----------------------------|------------------|----------|-----------------|--|--|--|
| T (MILE : 1 | | I Alfalfa I I 1 1 | /2 to 2 1/2 Town/ | ۸ | | | | | | |
| Amount of Mulch Required: None None | | lay, Alfalfa Hay - 1 1 | /2 to 2 1/2 1 ons// | Acre | | | | | | |
| Minimum Rate (lbs): | 3,000 | OR Tons: | 1.50 | Maximum Rate (lbs): | 5.000 OR | Tons: | 2.5 | | | |
| Willimum Rate (ibs). | 3,000 | OK TOILS. | 1.50 | iviaximum rate (ibs). | 5,000 OK | 10115. | 2.5 | | | |
| Additional Information: n/a | | | | | | | | | | |
| Additional information. 174 | | | | | | | | | | |
| | | | | | | | | | | |
| Applied | | | | | | | | | | |
| Type of Mulch Used: | | | | | | | | | | |
| Number of Bales: | | Bale Weight, lbs: | | Mulch Applied, Tons: | | Exacc/S | Shortage, Tons: | | | |
| Nulliber of Bales. | | bale Weight, ibs. | | Mulcii Applieu, Toris. | | EXCess/C | ononage, rons. | | | |
| Based on the information in this job sheet, all requirements for this seeding should be completed by: Year: | | | | | | | | | | |
| Was site prep completed as | | uno occumy oriouta | bo completed by. | | | | | | | |
| Seeding was completed on | | cording to the above | requirements | | | | | | | |
| ceeding was completed on | (Date) | ording to the above | requirements. | | | | | | | |
| | (, | | | | | | | | | |
| Producer's or Contr | actor's Signature | | | Date | | | | | | |
| For NRCS Use Only: | actor o Cignaturo | | | 24.0 | | | | | | |
| • | Does this seeding meet all NRCS standards and specifications and was implemented as planned above? | | | | | | | | | |
| Certified by: | | | | | | | | | | |
| · · · · · · · · · · · · · · · · · · · | | Date | Field Office | | | | | | | |
| NRCS Representative Date Field Office Additional Comments: | | | | | | | | | | |
| | | | | | | | | | | |
| When seeding is completed, return seeding plan to the Natural Resources Conservation Service with all receipts pertaining to this seeding (seed, fertilzer, lime, mulch, companion crop, labor, drill rental, etc.) and seed tests, including tags for yellow tag seed. | | | | | | | | | | |
| | Companion | orop, labor, urill feri | iai, eic.) anu seed | i tests, including tags for | yellow lay seed. | | | | | |