| | Job Sheet for Seeding |
|--|---|
| | |
| Planned by: | MO Mix 39 CP38E-42 Sand Date: County: |
| Client: | Address: |
| Farm(s): | Tract(s): Fields: CIN#: |
| Acres: | 1.00 Program: Wildlife Hebitet Planting (420) |
| Primary / Cost-shared FOTG Pra | actice: Wildlife Habitat Planting (420) Secondary FOTG Practice: |
| Primary Purpose of Seeding: | OR Ecological Site: |
| Site Soil Group: Additional Information: | OR Ecological Site. |
| Additional information. | |
| | Seedbed Preparation for Grass and/or Forb/Legume Seeding |
| Current Conditions: | |
| Conditions Desired at Planting: | Row crop stubble or tilled seed bed (to incorporate amendments, level field, or following construction operations) Killed sod (renovation; for example from introduced cool season grasses to native warm season grasses or from warm season grasses to a pollinator forb only planting) Suppressed existing vegetation, as for an inter-seeding of legumes on pastureland or native forbs on natural communities Slopes that are 4:1 or steeper that will receive a critical area planting (broadcast planting method only) |
| Seed Mix Contains: | Cool Season Grasses (majority) Warm Season Grasses (majority) Native Forbs Native/Introduced Legumes |
| Planting Method: | Interseeding: No Planting Season: |
| Acceptable Planting Dates: | Optimal Planting Dates: |
| | |
| Seedbed Prep: | |
| | |
| | |
| Seed Placement: | |
| | |
| Additional Instructions: | |
| Fertilizer and Lime | |
| Planned | |
| Is Fertilizer and/or Lime Needed? | |
| Program rules and/or plann | ned practice(s) do not require soil amendments (example - general CRP plantings, wildlife plantings on cropland) |
| Program rules and/or plann | ed practices require soil amendments be applied according to a recent soil test (less than 4 years old and after the most recent application of liming material |
| and/or fertilizer). Apply fert | tilizer and/or lime according to soil test recommendations as indicated below. led practices require soil amendments. Blanket rates will be used for a Critical Area Planting (342) as indicated below because a recent soil test is not available |
| Soil Test Recommendations | |
| Salt pH from soil test results: | Lime* N P_2O_5 K_2O |
| Can pri nom con toct recane. | lbs/ac |
| | Total Needed Not req. 0 0 0 |
| Blanket rates for constructed fills | when a lbs/ac #N/A |
| current soil test is not availab | ole: Total Needed #N/A |
| What legumes are included in the | e seed mix? None Alfalfa Cool Season Legumes other than Alfalfa Warm Season Legumes |
| When to apply lime: | |
| When to apply fertilizer: | No fertilizer will be applied. Soil nutrients are within ranges required for the planned species that will be planted. |

U.S. Department of Agriculture Natural Resources Conservation Service

Revised 9/2017

| | • | | | | _ | |
|---|--------------------|----------------|-------------------|----------------------|-------------------------|--|
| Additional Information: | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | _ . | |
| | | | | | | |
| Companion Crop Specifications | | | | | | |
| A companion crop may be establis The companion crop will be select | | | | nd/or legumes. | | |
| Planned Companion Crop: | None Planned Redto | op Spring Oats | Winter Wheat | Rate per Acre (lbs): | Total Bulk Lbs to Seed: | |
| | | | | | | |
| Additional Information: | | | | | | |
| | | | | | | |
| | | | | | | |
| Applied | | | | | | |
| | | | | | | |
| Companion Crop Planted: | | | To | tal Lbs Lbs/Ac | | |
| Redtop Spring | Oats Winter Wheat | | Bulk Lbs Planted: | 0.0 | | |

Revised 9/2017

| Seeding Mix Summary | | | | | | | | | | | | |
|---|---|---------------|-------------------------|--------------------|-------------|--------|-------------------------------|--------------------|---------------------|---------------|--|-----------------------------------|
| Acres to be Seeded: | 1.00 | | | | | | | | | | | |
| Practice Adjustment Factor: | | | Broadca | st Adjustme | ent Factor: | | - | Intersee | ding Adjust | tment Factor: | 1 | |
| Apply Adjustment Factors to: | | | | | | | | | | | | |
| Planned Seed Mix Seed Mix Purchased and Planted | | | | | | | | | | | | |
| | | | | | | | | | | | 0 111 00 | |
| Species | Acceptable Cultivar(s) | PLS lbs/ac | Total PLS lbs Needed | Bulk lbs Seeded | % Purity | % Germ | % Dormant/ Hard Seed | PLS lbs Planted | Shortage/ Excess | % of Mix | Cultivar OR Origin for MO Source Identified Class Seed | Meets all Require- ments |
| Splitbeard Bluestem, Andropogon ternarius | No cultivars available | 1.345 | 1.345 | | | | | | -1.345 | | | |
| Black-eyed Susan (B), Rudbeckia hirta | | 0.030 | 0.030 | | | | | | -0.030 | | | |
| Blue Lobelia, Lobelia siphilitica | | 0.008 | 0.008 | | | | | | -0.008 | | | |
| Canada Milkvetch, Astragalus canadensis | | 0.075 | 0.075 | | | | | | -0.075 | | | |
| Common Evening Primrose (B), Oenothera biennis | | 0.020 | 0.020 | | | | | | -0.020 | | | |
| False Sunflower, Heliopsis helianthoides | | 0.140 | 0.140 | | | | | | -0.140 | | | |
| Foxglove Beardtongue, Penstemon digitalis | | 0.030 | 0.030 | | | | | | -0.030 | | | |
| Golden Alexanders, Zizia aurea | | 0.075 | 0.075 | | | | | | -0.075 | | | |
| Gray Goldenrod, Solidago nemoralis | | 0.030 | 0.030 | | | | | | -0.030 | | | |
| Gray-headed Coneflower, Ratibida pinnata | Available: Sunglow (statewide) | 0.148 | 0.148 | | | | | | -0.148 | | | |
| Illinois Bundleflower, Desmanthus illinoensis | Available: Reno Germplasm (statewide) | 0.150 | 0.150 | | | | | | -0.150 | | | |
| Lanceleaf Coreopsis, Coreopsis lanceolata | | 0.157 | 0.157 | | | | | | -0.157 | | | |
| Maximilian Sunflower, Helianthus maximiliani | | 0.010 | 0.010 | | | | | | -0.010 | | | |
| Pale Purple Coneflower, Echinacea pallida | | 0.030 | 0.030 | | | | | | -0.030 | | | |
| Partridge Pea (A), Chamaecrista fasciculata | Available: Riley (statewide), Lark Selection (statewide) | 0.130 | 0.130 | | | | | | -0.130 | | | |
| Prairie Cinquefoil, Drymocallis arguta | | 0.019 | 0.019 | | | | | | -0.019 | | | |
| Purple Coneflower, Echinacea purpurea | | 0.360 | 0.360 | | | | | | -0.360 | | | |
| Purple Prairie Clover, Dalea purpurea | Available: Kaneb (statewide) | 0.353 | 0.353 | | | | | | -0.353 | | | |
| Rattlesnake Master, Eryngium yuccifolium | | 0.020 | 0.020 | | | | | | -0.020 | | | |
| Roundhead Lespedeza, Lespedeza capitata | Available: Kanoka (statewide) | 0.020 | 0.020 | | | | | | -0.020 | | | |

| | Available: Alexander Germplasm (statewide) | 0.040 | 0.040 | | | -0.040 | | |
|---|---|-------|-------|--|--|--------|--|--|
| Smooth Aster, Symphyotrichum laeve | | 0.020 | 0.020 | | | -0.020 | | |
| Stiff Goldenrod, Oligoneuron rigidum, Solidago rigida | | 0.010 | 0.010 | | | -0.010 | | |
| Upright Prairie Coneflower, Ratibida columnifera | | 0.110 | 0.110 | | | -0.110 | | |
| White Prairie Clover, Dalea candida | | 0.097 | 0.097 | | | -0.097 | | |
| Wild Bergamot, Monarda fistulosa | | 0.068 | 0.068 | | | -0.068 | | |
| Yarrow, Achillea millefolium | | 0.008 | 0.008 | | | -0.008 | | |

| | Total Planned: | 3.503 | 3.503 | | | |
|---|----------------|-------|-------|--------------------------------------|--|--|
| Planned Grasses PLS/ft ² : | 6.67 | | | Total PLS lbs. Forbs Planted: | | |
| Planned Forbs/Legumes PLS/ft ² : | 20.00 | | | Total % of Mix: | | |
| Planned Total PLS/ft ² : | 26.67 | | | % of mix of annual/biennial species: | | |

Customer and Seed Dealer Please Note:

Only viable, high quality and adapted seed will be used. All seed shall have a current seed test within 10 months of the planting date that lists germination, purity, and hard seed as a percentage for determining pure live seed and lists the percent of weed seed present that meets State seed quality law standards. Seed test results must be provided to the customer and USDA for confirmation that seed planted meets specifications outlined in this plan. Seed must be clean and relatively free of weed seed and contaminants. Seed that has become wet, moldy, or otherwise damaged in transit or storage is not acceptable. Certified seed is preferred but not required unless Missouri Source Identified Class Seed, or yellow tag, is denoted as being required.

Legume seed shall be inoculated with the proper, viable Rhizobium bacteria species prior to planting according to Missouri Agronomy Technical Note MO-36, "Legume Inoculation." Pre-inoculated seed shall be planted prior to the expiration date on the inoculum tag or be re-inoculated with the appropriate inoculum within 24 hours prior to seeding. Inoculated seed will not be exposed to direct sunlight. Inoculation of legumes may be waived when a current pasture inventory (JS-AGRON-24 or similar inventory) shows that the legume species to be seeded occupies more than 5 percent of the vegetative cover on the site prior to seeding.

For "Acceptable Cultivar(s)," north Missouri (MO) is all counties north of Bates, Henry, Benton, Morgan, Moniteau, Cole, Osage, Gasconade, Franklin, and St. Louis Counties. South Missouri is all counties including and south of those listed. Refer to Native Cultivars or Selections Information Sheet (IS-MO723N) for guidance on selecting the appropriate native grass cultivars to meet objectives. In all cases where a cultivar(s) is/are noted under the Acceptable Cultivar(s) heading above, Missouri Source Identified Class Seed (yellow tag seed) may be used instead of a cultivar variety or selection.

The following information applies only if "Locally adapted Missouri Source Identified Class Seed (yellow tag) required" is shown under the "Acceptable Cultivar(s)" heading above: Locally adapted Missouri Source Identified Class Seed, or yellow tag, is defined as a native plant that source genetically originated in Missouri, was not introduced, and existed within the state borders prior to arrival of settlers. All seed from herbaceous material shall comply with Missouri seed laws including Missouri Crop Improvement Association guidance. All seed will comply with AOSCA (Association of Official Seed Certifying Agencies) certification procedures (including appropriate tagging) to include third-party verification by the Missouri Crop Improvement Association of source, genetic identity, and genetic purity of wildland collected or field or nursery grown plant germplasm materials. Seed must be Missouri origin (grown in Missouri) and certified as Missouri Source Identified Class. Source Identified Certification means: parent seed is collected from natural remnant Missouri populations; no selection, testing, or breeding for specific traits; production fields are inspected to verify species, source, and lack of noxious weeds; and seed is certified for purity and germination. If Missouri origin (grown) source identified class seed is not available, Missouri source identified class seed may be obtained from adjoining states.

| Mulching Specifications (After Grass Se | edina) | | | | | | | | | |
|--|-------------------------|-----------------------|-----------------|--------------------------------|-----------|-------|--------------------|-------------------|--|--|
| T (MILE : 1 = | | LL. AIG-IG- LL- | 1 1 /2 / 2 1 /2 | T /A | | | | | | |
| Amount of Mulch Required: None None None | | Hay, Alfalfa Hay - | 1 1/2 to 2 1/2 | : Tons/Acre | | | | | | |
| Minimum Rate (lbs): | 3,000 | OR Tor | s: 1.50 | Maximum Rate (lb | c): 5.000 | OR | Tons: | 2.5 | | |
| William Rate (ibs). | 3,000 | OK 101 | 1.50 | iviaximum rate (ib | 5). 5,000 | OK | 10115. | 2.5 | | |
| Additional Information: n/a | | | | | | | | | | |
| Additional information: 11/a | | | | | | | | | | |
| | | | | | | | | | | |
| Applied | | | | | | | | | | |
| Type of Mulch Used: | | | | | | | | | | |
| Number of Bales: | | Bale Weight, lb | .c. | Mulch Applied, Tor | 201 | | Evenes/Sho | ortage, Tons: | | |
| Number of Bales. | | bale Weight, ic | 5. | ividicii Applied, Toi | 15. | | EXC622/2110 | ntage, rons. | | |
| Based on the information in this job she | et all requirements for | r this seeding sho | uld he comple | ated by: | | Year: | | | | |
| Was site prep completed a | | i tillo occurrig oric | ala bo comple | | | 1001. | | | | |
| Seeding was completed on | | cording to the ab | ove requireme | ante | | | | | | |
| occuring was completed on | (Date) | cording to the ab | ovo roquironio | ono. | | | | | | |
| | (, | | | | | | | | | |
| Producer's or Cont | ractor's Signature | | | Date | | | | | | |
| For NRCS Use Only: | actor o Cignaturo | | | 24.0 | | | | | | |
| Does this seeding meet all NRCS standar | ds and specifications | and was impleme | ented as plann | ed above? | | | | | | |
| Certified by: | | | | | | | | | | |
| · · · · · · · · · · · · · · · · · · · | Date | | Fie | eld Office | | | | | | |
| · | | | | | | | | | | |
| | Additional Comments: | | | | | | | | | |
| When seeding is completed | | | | ervation Service with all rece | | | ing (seed, fertilz | zer, lime, mulch, | | |
| companion crop, labor, drill rental, etc.) and seed tests, including tags for yellow tag seed. | | | | | | | | | | |