

### Job Sheet for Seeding

Planned by: MO Mix 40 Date: \_\_\_\_\_ County: \_\_\_\_\_  
Client: \_\_\_\_\_ Address: \_\_\_\_\_  
Farm(s): \_\_\_\_\_ Tract(s): \_\_\_\_\_ Fields: \_\_\_\_\_ CIN#: \_\_\_\_\_  
Acres: 1.00 Program: \_\_\_\_\_  
Primary / Cost-shared FOTG Practice: Wildlife Habitat Planting (420) Secondary FOTG Practice: \_\_\_\_\_  
Primary Purpose of Seeding: \_\_\_\_\_  
Site Soil Group: \_\_\_\_\_ OR Ecological Site: \_\_\_\_\_  
Additional Information: \_\_\_\_\_

### Seedbed Preparation for Grass and/or Forb/Legume Seeding

Current Conditions: \_\_\_\_\_

Conditions Desired at Planting:

- Row crop stubble or tilled seed bed (to incorporate amendments, level field, or following construction operations)
- Killed sod (renovation; for example from introduced cool season grasses to native warm season grasses or from warm season grasses to a pollinator forb only planting)
- Suppressed existing vegetation, as for an inter-seeding of legumes on pastureland or native forbs on natural communities
- Slopes that are 4:1 or steeper that will receive a critical area planting (broadcast planting method only)

Seed Mix Contains:  Cool Season Grasses (majority)  Warm Season Grasses (majority)  Native Forbs  Native/Introduced Legumes

Planting Method: \_\_\_\_\_ Interseeding: No Planting Season: \_\_\_\_\_  
Acceptable Planting Dates: \_\_\_\_\_ Optimal Planting Dates: \_\_\_\_\_

Seedbed Prep: \_\_\_\_\_

Seed Placement: \_\_\_\_\_

Additional Instructions: \_\_\_\_\_

### Fertilizer and Lime

**Planned**

Is Fertilizer and/or Lime Needed?

- Program rules and/or planned practice(s) do not require soil amendments (example - general CRP plantings, wildlife plantings on cropland)
- Program rules and/or planned practices require soil amendments be applied according to a recent soil test (less than 4 years old and after the most recent application of liming material and/or fertilizer). Apply fertilizer and/or lime according to soil test recommendations as indicated below.
- Program rules and/or planned practices require soil amendments. Blanket rates will be used for a Critical Area Planting (342) as indicated below because a recent soil test is not available (as for constructed fills and finished grades).

**Soil Test Recommendations**

Salt pH from soil test results: \_\_\_\_\_

	Lime*	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
lbs/ac				
Total Needed	Not req.	0	0	0

Blanket rates for constructed fills when a current soil test is not available:

	Lime*	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
lbs/ac	#N/A			
Total Needed	#N/A			

What legumes are included in the seed mix?  None  Alfalfa  Cool Season Legumes other than Alfalfa  Warm Season Legumes

When to apply lime: \_\_\_\_\_

When to apply fertilizer: \_\_\_\_\_

No fertilizer will be applied. Soil nutrients are within ranges required for the planned species that will be planted.

Additional Information:

---

---

---

---

---

**Companion Crop Specifications**

A companion crop may be established prior to or at the same time as a permanent seeding of grass, forbs, and/or legumes. The companion crop will be selected and managed to minimize competition with the permanent seeding.

Planned Companion Crop:  None Planned  Redtop  Spring Oats  Winter Wheat      Rate per Acre (lbs): \_\_\_\_\_      Total Bulk Lbs to Seed: \_\_\_\_\_

Additional Information:

---

**Applied**

Companion Crop Planted:

Redtop  Spring Oats  Winter Wheat

Bulk Lbs Planted:

Total Lbs	Lbs/Ac
	0.0

Seeding Mix Summary														
Acres to be Seeded:		1.00												
Practice Adjustment Factor:		Broadcast Adjustment Factor:			Interseeding Adjustment Factor:								1	
Apply Adjustment Factors to:														
Planned Seed Mix				Seed Mix Purchased and Planted										
Species	Acceptable Cultivar(s)	PLS lbs/ac	Total PLS lbs Needed	Bulk lbs Seeded	% Purity	% Germ	% Dormant/ Hard Seed	PLS lbs Planted	Shortage/ Excess	% of Mix	Cultivar OR Origin for MO Source Identified Class Seed	Meets all Require- ments		
Little Bluestem, <i>Schizachyrium scoparium</i>	Aldous (statewide), Camper (north MO), Cimarron (south MO - sandy sites only), Ozark (statewide), Yellow Tag (statewide)	2.920	2.920						-2.920					
Sideoats Grama, <i>Bouteloua curtipendula</i>	Butte (north MO), El Reno (statewide), Trailway (statewide), Yellow Tag (statewide)	2.200	2.200						-2.200					
Black-eyed Susan (B), <i>Rudbeckia hirta</i>		0.075	0.075						-0.075					
Canada Milkvech, <i>Astragalus canadensis</i>		0.020	0.020						-0.020					
Common Evening Primrose (B), <i>Oenothera biennis</i>		0.030	0.030						-0.030					
Common Milkweed, <i>Asclepias syriaca</i>		0.220	0.220						-0.220					
False Sunflower, <i>Heliopsis helianthoides</i>		0.050	0.050						-0.050					
Foxglove Beardtongue, <i>Penstemon digitalis</i>		0.019	0.019						-0.019					
Golden Alexanders, <i>Zizia aurea</i>		0.070	0.070						-0.070					
Gray Goldenrod, <i>Solidago nemoralis</i>		0.010	0.010						-0.010					
Gray-headed Coneflower, <i>Ratibida pinnata</i>	Available: Sunglow (statewide)	0.055	0.055						-0.055					
Illinois Bundleflower, <i>Desmanthus illinoensis</i>	Available: Reno Germplasm (statewide)	0.150	0.150						-0.150					
Lanceleaf Coreopsis, <i>Coreopsis lanceolata</i>		0.100	0.100						-0.100					
Maximilian Sunflower, <i>Helianthus maximiliani</i>		0.010	0.010						-0.010					

Pale Purple Coneflower, <i>Echinacea pallida</i>		0.160	0.160							-0.160			
Partridge Pea (A), <i>Chamaecrista fasciculata</i>	Available: Riley (statewide), Lark Selection (statewide)	0.100	0.100							-0.100			
Prairie Cinquefoil, <i>Drymocallis arguta</i>		0.030	0.030							-0.030			
Purple Prairie Clover, <i>Dalea purpurea</i>	Available: Kaneb (statewide)	0.100	0.100							-0.100			
Rattlesnake Master, <i>Eryngium yuccifolium</i>		0.020	0.020							-0.020			
Roundhead Lespedeza, <i>Lespedeza capitata</i>	Available: Kanoka (statewide)	0.020	0.020							-0.020			
Showy Ticktrefoil, <i>Desmodium canadense</i>	Available: Alexander Germplasm (statewide)	0.040	0.040							-0.040			
Skyblue Aster, <i>Symphyotrichum oolentangiense</i>		0.010	0.010							-0.010			
Stiff Goldenrod, <i>Oligoneuron rigidum</i> , <i>Solidago rigida</i>		0.010	0.010							-0.010			
Swamp Milkweed, <i>Asclepias incarnata</i>		0.080	0.080							-0.080			
Upright Prairie Coneflower, <i>Ratibida columnifera</i>		0.100	0.100							-0.100			
White Prairie Clover, <i>Dalea candida</i>		0.029	0.029							-0.029			
Wild Bergamot, <i>Monarda fistulosa</i>		0.070	0.070							-0.070			
<b>Total Planned:</b>		<b>6.698</b>	<b>6.698</b>										
Planned Grasses PLS/ft <sup>2</sup> :		<u>24.17</u>											
Planned Forbs/Legumes PLS/ft <sup>2</sup> :		<u>16.00</u>											
Planned Total PLS/ft <sup>2</sup> :		<u>40.17</u>											
			Total PLS lbs. Forbs Planted: _____										
			Total % of Mix: _____										
			% of mix of annual/biennial species: _____										
<b>Customer and Seed Dealer Please Note:</b>													

Only viable, high quality and adapted seed will be used. All seed shall have a current seed test within 10 months of the planting date that lists germination, purity, and hard seed as a percentage for determining pure live seed and lists the percent of weed seed present that meets State seed quality law standards. Seed test results must be provided to the customer and USDA for confirmation that seed planted meets specifications outlined in this plan. Seed must be clean and relatively free of weed seed and contaminants. Seed that has become wet, moldy, or otherwise damaged in transit or storage is not acceptable. Certified seed is preferred but not required unless Missouri Source Identified Class Seed, or yellow tag, is denoted as being required.

Legume seed shall be inoculated with the proper, viable Rhizobium bacteria species prior to planting according to Missouri Agronomy Technical Note MO-36, "Legume Inoculation." Pre-inoculated seed shall be planted prior to the expiration date on the inoculum tag or be re-inoculated with the appropriate inoculum within 24 hours prior to seeding. Inoculated seed will not be exposed to direct sunlight. Inoculation of legumes may be waived when a current pasture inventory (JS-AGRON-24 or similar inventory) shows that the legume species to be seeded occupies more than 5 percent of the vegetative cover on the site prior to seeding.

For "Acceptable Cultivar(s)," north Missouri (MO) is all counties north of Bates, Henry, Benton, Morgan, Moniteau, Cole, Osage, Gasconade, Franklin, and St. Louis Counties. South Missouri is all counties including and south of those listed. Refer to Native Cultivars or Selections Information Sheet (IS-MO723N) for guidance on selecting the appropriate native grass cultivars to meet objectives. In all cases where a cultivar(s) is/are noted under the Acceptable Cultivar(s) heading above, Missouri Source Identified Class Seed (yellow tag seed) may be used instead of a cultivar variety or selection.

**The following information applies only if "Locally adapted Missouri Source Identified Class Seed (yellow tag) required" is shown under the "Acceptable Cultivar(s)" heading above:** Locally adapted Missouri Source Identified Class Seed, or yellow tag, is defined as a native plant that source genetically originated in Missouri, was not introduced, and existed within the state borders prior to arrival of settlers. All seed from herbaceous material shall comply with Missouri seed laws including Missouri Crop Improvement Association guidance. All seed will comply with AOSCA (Association of Official Seed Certifying Agencies) certification procedures (including appropriate tagging) to include third-party verification by the Missouri Crop Improvement Association of source, genetic identity, and genetic purity of wildland collected or field or nursery grown plant germplasm materials. Seed must be Missouri origin (grown in Missouri) and certified as Missouri Source Identified Class. Source Identified Certification means: parent seed is collected from natural remnant Missouri populations; no selection, testing, or breeding for specific traits; production fields are inspected to verify species, source, and lack of noxious weeds; and seed is certified for purity and germination. If Missouri origin (grown) source identified class seed is not available, Missouri source identified class seed may be obtained from adjoining states.

**Mulching Specifications (After Grass Seeding)**

Type of Mulch Required:  None  Straw, Grass Hay, Alfalfa Hay - 1 1/2 to 2 1/2 Tons/Acre

Amount of Mulch Required for This Site:  
 Minimum Rate (lbs): 3,000 OR Tons: 1.50 Maximum Rate (lbs): 5,000 OR Tons: 2.5

Additional Information: n/a

**Applied**

Type of Mulch Used: \_\_\_\_\_  
 Number of Bales: \_\_\_\_\_ Bale Weight, lbs: \_\_\_\_\_ Mulch Applied, Tons: \_\_\_\_\_ Excess/Shortage, Tons: \_\_\_\_\_

Based on the information in this job sheet, all requirements for this seeding should be completed by: \_\_\_\_\_ Year: \_\_\_\_\_  
 Was site prep completed as outlined above? \_\_\_\_\_  
 Seeding was completed on \_\_\_\_\_ according to the above requirements.  
 (Date)

\_\_\_\_\_  
 Producer's or Contractor's Signature Date

**For NRCS Use Only:**

Does this seeding meet all NRCS standards and specifications and was implemented as planned above? \_\_\_\_\_

Certified by: \_\_\_\_\_  
 NRCS Representative Date Field Office

Additional Comments: \_\_\_\_\_

When seeding is completed, return seeding plan to the Natural Resources Conservation Service with all receipts pertaining to this seeding (seed, fertilizer, lime, mulch, companion crop, labor, drill rental, etc.) and seed tests, including tags for yellow tag seed.